


Colorectal Surgical Pathway

Patient Journey

Straight to Surgery



Surgery

4-6 weeks post-surgery

Rehak

Handoff from hospital to GM Active contact. GM Active contact participant 4 weeks post-surgery via telephone call

4-6 weeks post-surgery:

Assessment clinic 4-6 weeks and 18 weeks post-surgery
12 week free gym membership
Personalised exercise programme
Graded approach to resuming exercise post-treatment
Focus on behavioral change and long-term adoption of exercise plan















4-6 weeks post-surgery:

Assessment clinic 4-6 weeks and 18 weeks post-surgery
12 week free gym membership
Personalised exercise programme
Graded approach to resuming exercise post-treatment
Focus on behavioral change and long-term adoption of exercise plan















utcome

30 day mortality 90 day mortality Clavien Dindo POMs day 7 and 15 30 day readmission 2 year survival Patent

EQ5D WHO-DAS 2.0 FACT FRAILTY

PHYSIOLGICAL

Exclusions: Jnstable cardiac or resp disease (recent MI) Syncope attacks SaO <85% at rest PE < 2 weeks Patients unable to maintain a steady walking pag

Other potential contraindications:
Severe hypertension
Uncontrolled arrhythmia
Symptomatic Severe aortic stenosis
AAA > 8cms
Advanced pregnancy
Known HOCM

CPET

Unstable cardiac or resp disease (recent MI)
Syncope attacks
PE < 2 weeks
Patients unable to cycle











